

Indications for Criticality at Zero Curvature in a 4d Regge Model of Euclidean Quantum Gravity*

Wolfgang Beirl^a and Bernd A. Berg^{b,c}

^aMarket Research Inc., Paradise Island, Bahamas

^bDepartment of Physics, The Florida State University, Tallahassee, FL 32306, USA

^cSchool of Computational Science and Information Technology,
The Florida State University, Tallahassee, FL 32306, USA

We re-examine the approach to four-dimensional Euclidean quantum gravity based on the Regge calculus. A cut-off on the link lengths is introduced and consequently the gravitational coupling and the cosmological constant become independent parameters. We determine the zero curvature, $\langle R \rangle = 0$, line in the coupling constant plane by numerical simulations. When crossing this line we find a strong, probably first order, phase transition line with indications of a second order endpoint. Beyond the endpoint the transition through the $\langle R \rangle = 0$ line appears to be a crossover. Previous investigations, using the Regge or the Dynamical Triangulation approach, dealt with a limit in which the first order transition prevails.

1. Introduction

Euclidean quantum gravity in four dimensions is formally defined by the path integral

$$Z = \int Dg e^{-S_{EH}[g]}. \quad (1)$$

The Einstein-Hilbert action $S_{EH}[g]$ is a function of the 4-geometry g , $S_{EH}[g] = -L_P^{-2}R[g] + \Lambda V[g]$. The Planck length L_P determines the gravitational coupling constant for the total curvature $R[g]$ and the total 4-volume $V[g]$ enters the action if the cosmological constant Λ is non-zero.

In order to *define* the integral (1), we consider a lattice approximation, so that the path-integral becomes the limit of a series of well-defined and finite integrals. A direct route for such an approximation is provided by the Regge calculus [1], which is suitable for simulations [2,3]. One considers a simplicial lattice of fixed connectivity, which subdivides a 4-manifold of given topology in N_4 4-simplices, N_3 tetraeder, N_2 triangles, N_1 links and N_0 vertices. As we assign a length to each link, a 4-geometry is defined in the Euclidean sector, if the triangle inequalities are fulfilled

for every (sub-)simplex. In the following we will use the squared lengths q_l associated with each link l as the basic variables and a given set of such variables defines a 4-geometry $[q]$ for a given simplicial lattice. Variation of the variables q_l allows us to approximate the path integral (1). One can calculate the Regge action $R[q] = \sum_t A_t \delta_t$ as a sum over all triangles t with area A_t multiplied by their associated deficit angles δ_t . The 4-volume is calculated as a sum over the volume of all 4-simplices s , $V[q] = \sum_s V_s$. We then approximate the gravitational action by $S_{EH} \rightarrow S[q] = -L_P^{-2}R[q] + \Lambda V[q]$. It has been demonstrated how the geometrical variables of the Regge calculus converge towards the corresponding variables of General Relativity in the classical continuum limit [1,4]. It requires an upper bound on the link lengths, $q_l < Q$, and a limit F on the 'fatness' $\phi_s = V_s / \max(q_l^2)$ of each 4-simplex, $\phi_s > F$. The continuum limit is then reached for $N_0 \rightarrow \infty$, $Q \rightarrow 0$ with $F > 0$ (Cheeger et al. [4]).

In order to replace the path integral (1) by a finite integral on the simplicial lattice,

$$Z = \int Dq e^{-S[q]}. \quad (2)$$

*This work was in part supported by the US Department of Energy under contract DE-FG02-97ER41022.

we need to define the integration measure. A simple choice is the uniform measure

$$Dq = \int_0^Q \prod_l dq_l \Theta_F[q] \quad (3)$$

where the function Θ_F equals 1 when all triangle inequalities are fulfilled and $\phi_s > F$ for all 4-simplices; it is 0 otherwise. The integration limit Q is necessary for a well-defined integral, but it has no physical meaning in itself. Indeed, simple re-scaling $q_l \rightarrow Qq_l$ formally removes it and leads to the integral

$$Z(\beta, \lambda) = \int_0^1 \prod_l dq_l \Theta_F[q] e^{\beta R[q] - \lambda V[q]} \quad (4)$$

where $\beta = QL_P^{-2}$ and $\lambda = Q^2\Lambda$.

In previous numerical investigations [5,6] the upper limit Q was infinite which corresponds to the limit $\beta \rightarrow \infty$ and $\lambda \rightarrow \infty$ in the integral (4).

In the following we will numerically evaluate the integral (4) in order to determine the phase structure by varying the coupling parameters β and λ . Our results suggest a physically interesting continuum limit for finite values of the coupling parameters.

2. Results

The numerical computations have been performed for the integral

$$Z(\beta, \lambda) = \int_0^c \prod_l dq_l \Theta_F[q] e^{\beta R[q] - \lambda V[q]} \quad (5)$$

with $c = 10$ instead of (4) with $c = 1$. This means that the coupling constants and expectation values need to be adjusted accordingly, e.g. $\beta \rightarrow c\beta$, $\lambda \rightarrow c^2\lambda$ etc. We set $F = 10^{-6}$ but our results seem to be independent of the specific value for a relatively wide range $F = 10^{-8}$ to 10^{-5} . We used a hypercubic regular triangulation of the 4-torus [2,3], with $N_0 = 4^4, 4^3 \times 10, 6^4$ and 8^4 vertices.

From simulations with $\lambda = 1, 2, 3, 4$ and varying β we find expectation values $\langle R \rangle$, which are (almost) zero for $\beta_0(\lambda)$ values; β_0 initially decreases for increasing λ . Then, as depicted in figure 1, we find that β_0 increases for $\lambda = 4, 5, 6, 7$.




Figure 1. Hysteresis loops.




Figure 2. Function $\beta_0(\lambda)$ for which $\langle R \rangle = 0$ holds.

Furthermore, the transition from $\langle R \rangle < 0$ to $\langle R \rangle > 0$ is no longer defined clearly, because we have to deal with a significant hysteresis for large λ . The line $\beta_0(\lambda)$ is depicted in figure 2 for different lattice size. The curve $\beta_0(\lambda)$ is interesting, since a vanishing curvature is a desired ingredient for a physical continuum limit.

As we approach the turning point $\lambda_c \sim 4.0$ from smaller λ , the susceptibility $\chi_q = N_1^{-1} (\langle \bar{q}^2 \rangle - \langle \bar{q} \rangle^2)$, with $\bar{q} = N_1^{-1} \sum_l q_l$, increases significantly; this increase is more pronounced for larger lattice size as depicted in figure 3.

Next, we performed calculations varying λ at $\beta = 0.175$ and 0.172 for the lattice size $4^3 \times 10$. As expected, we find that the curvature $\langle R \rangle$ increases initially, exhibiting a transition from negative to positive values, but when λ increases further a second transition back to negative values is




Figure 3. Susceptibility χ_q as function of λ at β_0 for increasing lattice size.




Figure 4. Susceptibility χ_R as function of λ for two β values.

seen. We depict $\chi_R = N_2^{-1}(\langle \overline{R}^2 \rangle - \langle \overline{R} \rangle^2)$, measuring curvature fluctuations, in figure 4; for $\beta = 0.175$ one can distinguish two peaks corresponding to the two transitions. These peaks merge into one at $\beta = 0.172$.

We also measured the 2-point correlation function $K(d) = \langle q_0 q_d \rangle - \langle q_0 \rangle^2$ along the elongated direction on the $4^3 \times 10$ lattice for $\beta_0(\lambda)$ and found that the function $K(d)$ is significantly different from zero for $d > 1$. Negative values alternate with positive values, which indicates a non-trivial interaction. The correlations increase for $\lambda \rightarrow \lambda_c$.

Although we are not able to provide a complete finite size scaling analysis, our data suggest the following: In the region $\lambda > \lambda_c$, with $\lambda \sim 4.0$, one deals with a strong phase transition, which is

likely of first order. In the region $\lambda < \lambda_c$ we are currently unable to distinguish between a cross-over transition or a higher-order phase transition. However the end-point λ_c itself is a natural candidate for a second-order phase transition. The increase of susceptibility χ_q and the link correlations $K(d)$ approaching λ_c from smaller values confirm this picture.

3. Summary and Conclusions

Our simulations differ from those of the previous literature by introducing the limit Q in the functional integral (3). While a simple re-scaling removes Q formally from the partition function (4), our model remains more general than those simulated in the literature, which are recovered in our limit of large λ . Remarkably, the new model appears to have a second order fixed point in the $\lambda - \beta$ plane, which deserves further study.

REFERENCES

1. T. Regge, Nuovo Cimento **9** (1961) 558.
2. B.A. Berg, Phys. Rev. Lett. **55** (1985) 904.
3. H.W. Hamber and R.M. Williams, Phys. Lett. B **157** (1985) 368.
4. R. Friedberg and T.D. Lee, Nucl. Phys. B **242** (1984) 145; G. Feinberg, R. Friedberg, T.D. Lee and H.C. Ren, Nucl. Phys. B **245** (1984) 343; H. Cheeger, W. Müller and R. Schrader, Commun. Math. Phys. **92** (1984) 405.
5. M. Gross and H. Hamber, Nucl. Phys. B **364** (1991) 703; C. Holm and W. Janke, Phys. Lett. B **335** (1994) 143.
6. W. Beirl, P. Homolka, B. Krishnan, M. Markum and J. Riedler, Nucl. Phys. B (Proc. Suppl.) **42**, 710 (1995); J. Riedler, W. Beirl, E. Bittner, A. Hauke, P. Homolka and H. Markum Class. Quant. Grav. **16**, 1163 (1999).